From Blanchard, Devaney, and Hall 3<sup>rd</sup> Edition:

## EXERCISES FOR APPENDIX B

In Exercises 1–4, use the guess-and-test method to find the power series expansion centered at t = 0 for the general solution up to degree four, that is, up to and including the  $t^4$  term. (You may find the general solution using other methods and then find the Taylor series centered at t = 0 to check your computation if you like.)

1. 
$$\frac{dy}{dt} = y$$
  
3.  $\frac{dy}{dt} = -2ty$   
2.  $\frac{dy}{d2} = -y+1$   
4.  $\frac{dy}{dt} = t^2y+1$ 

In Exercise 5–8, find the power series expansion for the general solution up to degree four, that is, up to and including the  $t^4$  term.

5. 
$$\frac{dy}{dt} = -y + e^{2t}$$
  
6. 
$$\frac{dy}{dt} = 2y + \sin t$$
  
7. 
$$\frac{d^2y}{dt^2} + 2y = \cos t$$
  
8. 
$$\frac{d^2y}{dt^2} + 5\frac{dy}{dt} + y = \sin 2t$$

9. Verify that  $y(t) = \tan t$  is a solution of

$$\frac{dy}{dt} = y^2 + 1$$

and compute a power series solution to find the terms up to degree six (up to and including the  $t^6$  term) of the Taylor series centered at t = 0 of tan t.

In Exercises 10–13, find the general solution up to degree six, that is, up to and including the  $t^6$  term.

**10.** 
$$\frac{d^2y}{dt^2} + 2y = 0$$
  
**11.**  $\frac{d^2y}{dt^2} + \frac{dy}{dt} + y = 0$   
**12.**  $\frac{d^2y}{dt^2} + \frac{dy}{dt} + t^2y = \cos t$   
**13.**  $\frac{d^2y}{dt^2} + t\frac{dy}{dt} + y = e^{-2t}$ 

## Hints and Answers for Appendix B

- **1.** The Taylor series centered at t = 0 for  $y(t) = ke^t$ .
- **3.** The Taylor series centered at t = 0 for  $y(t) = ke^{-t^2}$ .
- 5.  $y(t) = a_0 + (-a_0 + 1)t + (a_0/2 + 1/2)t^2 + (-a_0/6 + 1/2)t^3 + (a_0/24 + 5/24)t^4 + \dots$
- 7.  $y(t) = a_0 + a_1t + (1/2 a_0)t^2 (a_1/3)t^3 + (a_0/6 1/8)t^4 + \dots$
- 9.  $\tan t = t + t^3/3 + 2t^5/15 + \dots$
- **11.**  $y(t) = a_0 + a_1 t + (-a_0/2 a_1/2) t^2 + (a_0/6) t^3 + (a_1/24) t^4 + (-a_0/120 a_1/120) t^5 + (a_0/720) t^6 + \dots$
- **13.**  $y(t) = a_0 + a_1 t + (1/2 a_0/2) t^2 + (-1/3 a_1/3) t^3 + (1/24 + a_0/8) t^4 + (a_1/15)t^5 + (11/720 a_0/48) t^6 + \dots$

(a) 
$$a_2 = -\frac{\nu(\nu+1)}{2}a_0$$
,  
 $a_3 = \frac{2-\nu(\nu+1)}{2}a_1$ ,  
 $a_4 = -\frac{6-\nu(\nu+1)}{12}\frac{\nu(\nu+1)}{2}a_0$   
(b) *Hint:* Note that  $a_{2n}$  has  $a_0$  as a factor

15.

- (b) *Hint*: Note that  $a_{2n}$  has  $a_0$  as a factor and  $a_{2n+1}$  has  $a_1$  as a factor. Also note that if v = n is a positive integer, then  $a_{n+2} = 0$ .
- (c) *Hint*: Use the formulas from part (a).

(d) 
$$P_3(t) = t - \frac{5}{3}t^3$$
,  
 $P_4(t) = 1 - 10t^2 + \frac{35}{3}t^4$ ,  
 $P_5(t) = t - \frac{14}{3}t^3 + \frac{21}{5}t^5$ ,  
 $P_6(t) = 1 - 21t^2 + 63t^4 - \frac{231}{5}t^6$   
(e) *Hint*: Use linearity.

**17.** 
$$y(t_0) = t - t^2 + t^3/2 - t^4/6 + \dots$$