- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.

- 1. Prove Theorem 1.7.2(a) for the multiplicative case.
- 2. Prove Theorem 1.7.2(b) for the additive case.
- 3. Prove Theorem 1.7.2(e).
- 4. Prove Theorem 1.7.2(f).
- 5. Prove Theorem 1.7.2(g).
- 6. Prove Theorem 1.7.4(c).
- 7. Prove Theorem 1.7.4(d).
- 8. Prove that if $r \ge 1$ is a real number, then $r^2 \ge r$ and $\frac{1}{r^2} \le \frac{1}{r}$.
- 9. Prove that if a set *A* has a supremum, then sup *A* is unique.
- 10. Prove that parts (a) and (c) of Theorem 1.7.9 are equivalent.