

Fake Quiz 4 Calc 3 11/20/2009

This is a fake quiz, this is *only* a fake quiz. In the event of an actual quiz, you'd have been given fair warning. Repeat: This is *only* a fake quiz.

1. Compute $\int_C (x^2 + y^2) dx - x dy$ along the quarter circle from (1,0) to (0,1).

Integrate the long way to get $-1 - \pi/4$.

2. Evaluate $\int_C (\sin y \sinh x + \cos y \cosh x) dx + (\cos y \cosh x - \sin y \sinh x) dy$ where C is the line segment from (1,0) to $(2, \frac{\pi}{2})$.

Integrate using the Fundamental Theorem for Line Integrals (the potential function is $f = \sin y \cosh x + \cos y \sinh x$) to get $\cosh 2 - \sinh 1$.

3. Evaluate $\iint_S \mathbf{F} \cdot \mathbf{n} dS$, where $\mathbf{F}(x,y,z) = 4x \mathbf{i} - 3y \mathbf{j} + 7z \mathbf{k}$ and S is the surface of the cube bounded by the coordinate planes and the planes $x = 1$, $y = 1$, and $z = 1$.

Integrate using the Divergence Theorem to get 8.

4. Evaluate $\iint_S \mathbf{F} \cdot \mathbf{n} dS$, where $\mathbf{F}(x,y,z) = x \mathbf{i} + y \mathbf{j} + 2z \mathbf{k}$ and S is the portion of the cone $z^2 = x^2 + y^2$ between the planes $z = 1$ and $z = 2$, oriented upwards.

Integrate the long way to get $14\pi/3$.

5. Evaluate $\int_C (x^2 - y) dx + x dy$, where C is the circle $x^2 + y^2 = 4$ with counterclockwise orientation.

Use Green's Theorem to get 8π .

6. Evaluate $\iint_S \langle x^3, x^2y, xy \rangle \cdot d\mathbf{S}$, where S is the surface of the solid bounded by $z = 4 - x^2$, $y + z = 5$, $z = 0$, and $y = 0$.

Use the Divergence Theorem to get $4608/35$.

7. Compute $\int_C \mathbf{F} \cdot d\mathbf{r}$ where $\mathbf{F}(x,y,z) = y \mathbf{i} + z \mathbf{j} - x \mathbf{k}$ and C is the line segment from (1,1,1) to

$(-3, 2, 0)$.

Integrate the long way to get $-13/2$.

8. Compute $\int_C \left\langle \ln(1+y), -\frac{xy}{1+y} \right\rangle \cdot d\mathbf{r}$ where C is the triangle with vertices $(0,0)$, $(2,0)$, and $(0,4)$.

Use Green's Theorem to get -4 .

9. Evaluate $\int_{(0,1)}^{(\pi,-1)} y \sin x \, dx - \cos x \, dy$

Use the Fundamental Theorem for Line Integrals (the potential function is $f = -y \cos x$ to get 0.

10. Compute $\iint_S \mathbf{F} \cdot \mathbf{n} \, dS$, where $\mathbf{F}(x,y,z) = 2y \mathbf{j} + \mathbf{k}$ and S is the portion of the paraboloid $z = x^2 + y^2$ below the plane $z = 4$ with positive orientation.

Use the long way to get -12π .