

Each problem is worth 10 points. For full credit provide complete justification for your answers.

1. Evaluate $\int \sin^5 \theta d\theta$.

$$\int (1 - \cos^2 \theta)^2 \sin \theta d\theta$$

$$1 = \sin^2 \theta + \cos^2 \theta$$

$$\sin^2 \theta = 1 - \cos^2 \theta$$

$$-1 \int (1 - u^2)^2 \sin \theta \frac{du}{-\sin \theta}$$

$$u = \cos \theta$$

$$du = -\sin \theta d\theta$$

$$-1 \int (1 - u^2)(1 - u^2) du$$

$$d\theta = \frac{du}{-\sin \theta}$$

$$-1 \int 1 - 2u^2 + u^4 du$$

$$-1 \left(u - \frac{2}{3} u^3 + \frac{1}{5} u^5 \right) + C$$

Great!

$$= \boxed{-\cos \theta + \frac{2}{3} \cos^3 \theta - \frac{1}{5} \cos^5 \theta + C}$$

2. Evaluate $\int \ln(2x+1) dx$.

$$\text{let } u = 2x + 1$$

$$\frac{du}{dx} = 2, \quad \frac{du}{2} = dx$$

$$\int \ln(u) \frac{du}{2} \quad (\text{use line 100})$$

$$= (u \ln u - u) \frac{1}{2} + C$$

$$= \frac{1}{2} \left((2x+1) \ln|2x+1| - (2x+1) \right) + C$$

okay.

3. Evaluate $\int \frac{7}{(x+5)(x-2)} dx$. partial fractions

I wish $\frac{A}{(x+5)} + \frac{B}{(x-2)}$

$$7 = A(x-2) + B(x+5)$$

$x=2$ then $7 = B(7)$ so $B=1$

$x=-5$ then $7 = A(-7)$ so $A=-1$

Excellent!

$$\int \frac{-1}{(x+5)} dx + \int \frac{1}{(x-2)} dx = -\ln|x+5| + \ln|x-2| + C$$

4. Find the present value of an income stream of \$2000 per year, for a period of 10 years, if the continuous interest rate is 7%.

$$y = pe^{rt}$$

$$y = 2000e^{.07(x)}$$

$$u = .07x$$

$$\frac{du}{.07x} = dx$$

$$= \int_0^{10} 2000e^{.07x} dx = 2000 \int_0^{10} e^{.07x} dx$$

$$= 2000 \int_0^{10} e^u \frac{du}{.07}$$

$$= 2000 \left| \frac{1}{.07} e^{.07x} \right|_0^{10}$$

$$= 2000 \left(\frac{1}{.07} e^{.7} - \frac{1}{.07} \right)$$

Nice!

5. Find the median of the probability density function $p(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{1}{3}e^{-t/3} & \text{if } t \geq 0 \end{cases}$.

Median function

$$\frac{1}{2} = \int_{-\infty}^m p(t) dt$$

$$\frac{1}{2} = \int_{-\infty}^0 0 dt + \int_0^m \frac{1}{3} e^{-t/3} dt = \lim_{b \rightarrow \infty} \int_b^0 0 dt + \int_0^m \frac{1}{3} e^{-t/3} dt.$$

$$\frac{1}{2} = 0 + \left[-e^{-t/3} \right]_0^m$$

$$\frac{1}{2} = -e^{-m/3} + e^0$$

$$-\frac{1}{2} = -e^{-m/3}$$

$$e^{-m/3} = \frac{1}{2}$$

$$\frac{-m}{3} = \ln \frac{1}{2}$$

$$m = 3 \ln 2$$

Great



6. Find the length of the curve $y = \ln(1 - x^2)$ for $0 \leq x \leq \frac{1}{2}$.

$$y' = \frac{1}{1-x^2} \cdot -2x$$

$$\begin{aligned} \text{Arc Length} &= \int_a^b \sqrt{1 + [f'(x)]^2} dx \\ &= \int_0^{\frac{1}{2}} \sqrt{1 + \left(\frac{-2x}{1-x^2}\right)^2} dx \\ &= \int_0^{\frac{1}{2}} \sqrt{1 + \frac{4x^2}{1-2x^2+x^4}} dx \\ &= \int_0^{\frac{1}{2}} \sqrt{\frac{1+2x^2+x^4}{1-2x^2+x^4}} dx \\ &= \int_0^{\frac{1}{2}} \sqrt{\frac{(1+x^2)^2}{(1-x^2)^2}} dx \end{aligned}$$

$$\begin{aligned} &= \int_0^{\frac{1}{2}} \frac{1+x^2}{1-x^2} dx \\ &= \int_0^{\frac{1}{2}} \left(\frac{1-x^2}{1-x^2} + \frac{2}{1-x^2} \right) dx \\ &= \left[-x + 2 \cdot \frac{1}{2} \ln \left| \frac{x+1}{x-1} \right| \right]_0^{\frac{1}{2}} \\ &= \left(-\frac{1}{2} + \ln \left| \frac{3/2}{-1/2} \right| \right) - \left(0 - \ln \left| \frac{1}{-1} \right| \right) \\ &= -\frac{1}{2} + \ln 3 + \ln 1 \\ &= -\frac{1}{2} + \ln 3 \end{aligned}$$

7. Bunny is a Calculus student at Enormous State University, and she's having some trouble. Bunny says "Ohmygod, this is sooooo impossible. These integrals are crazy, and even once they said we could use the table it's still so confusing. There was this one on the test, and we were supposed to integrate tangent, right? So I found Line 75 on our table, it's the same book as you guys have, and I used it, and so there was 0 in the bottom of the fraction and I marked "Does Not Exist" for the answer, but they said it was wrong. Is that totally unfair or what?"

Explain clearly to Bunny what limitations the Table of Integrals might have that affected her, and how she might have used it better.

line 75 is about $\int \tan^n u \, du = \frac{1}{n-1} \tan^{n-1} u - \int \tan^{n-2} u \, du$
it looks like it can solve all problems of $\tan^n u$ with n can be any number. But that's not true.

When $n=1$, $\frac{1}{n-1}$ does not exist. It doesn't mean the integral doesn't exist, but just that this function (line 75) does not apply. If Bunny uses line 12 instead, which is

$$\int \tan u \, du = \ln|\sec u| + C, \text{ she'd get}$$

the correct answer.

When using the table, it is important to have basic calculus knowledge and know the conditions and limitations.

Excellent!

8. Evaluate $\int \tan^n \theta \sec^2 \theta d\theta$.

Since Sec is even

$$u = \tan \theta \quad d\theta = \frac{du}{\sec^2 \theta}$$

$$du = \sec^2 \theta d\theta$$

$$\underline{1 + \tan^2 \theta = \sec^2 \theta}$$

$$\int \tan^n \theta \sec^2 \theta d\theta$$

$$\int u^n \cancel{\sec^2 \theta} \cdot \frac{du}{\cancel{\sec^2 \theta}}$$

$$\int u^n \cdot du$$

Great!

$$\frac{1}{n+1} u^{n+1}$$

$$= \frac{1}{n+1} (\tan \theta)^{n+1} + C$$

$$\int \sqrt{a^2 - u^2} du = \frac{u}{2} \sqrt{a^2 - u^2} + \frac{a^2}{2} \sin^{-1} \frac{u}{a} + C$$

9. Derive Line 30 on the Table of Integrals.

Let $u = a \sin \theta$

$$\frac{du}{d\theta} = a \cos \theta$$

$$du = a \cos \theta d\theta$$

$$\sin \theta = \frac{u}{a}$$

$$\theta = \sin^{-1} \frac{u}{a}$$

$$\text{Left} = \int \sqrt{a^2 - u^2} du$$

$$= \int \sqrt{a^2 - a^2 \sin^2 \theta} \cdot a \cos \theta d\theta$$

$$= \int a \cos \theta \cdot a \cos \theta d\theta$$


$$= a^2 \int \cos^2 \theta d\theta$$

line 64 $= a^2 \left[\frac{1}{2} \theta + \frac{1}{4} \sin 2\theta + C \right]$

$$= \frac{a^2}{2} \sin^{-1} \frac{u}{a} + \frac{a^2}{4} 2 \sin \theta \cos \theta + C$$

$$= \frac{a^2}{2} \sin^{-1} \frac{u}{a} + \frac{a^2}{4} \cdot 2 \cdot \frac{u}{a} \cdot \frac{\sqrt{a^2 - u^2}}{a} + C$$

$$= \frac{a^2}{2} \sin^{-1} \frac{u}{a} + \frac{u}{2} \sqrt{a^2 - u^2} + C = \text{Right}$$



If do not use line 64 of table:

can use $\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$

to get the same result.

Excellent!

$$y' = \frac{2x}{w} = \frac{x}{5}$$

10. Jon wants to build the world's biggest barrel. It will be shaped like the region obtained by rotating the area below $y = x^2/10$ between $x = -20$ and $x = 20$ around the x -axis. Find the surface area of Jon's barrel.

$$\text{Surface Area} = \int_a^b 2\pi f(x) \sqrt{1 + [f'(x)]^2} dx$$

$$= \int_{-20}^{20} 2\pi \frac{x^2}{w} \sqrt{1 + \left(\frac{x}{5}\right)^2} dx$$

$$= \pi \int_{-20}^{20} \frac{x^2}{5} \sqrt{\frac{25 + x^2}{25}} dx$$

$$= \frac{\pi}{25} \int_{-20}^{20} x^2 \sqrt{25 + x^2} dx$$

$$= \frac{\pi}{25} \left[\frac{x}{8} (25 + 2x^2) \sqrt{25 + x^2} - \frac{625}{8} \ln(x + \sqrt{25 + x^2}) \right]_{-20}^{20}$$

$$= \frac{\pi}{25} \left[\frac{20}{8} (25 + 800) \sqrt{25 + 400} - \frac{625}{8} \ln(20 + \sqrt{25 + 400}) \right.$$

$$\left. - \left[\frac{-20}{8} (25 + 800) \sqrt{25 + 400} - \frac{625}{8} \ln(-20 + \sqrt{25 + 400}) \right] \right\}$$