

1. For any sets  $A, B,$  and  $C, A \cup (B \cap C) \subseteq A \cup B.$

Let  $x \in A \cup (B \cap C).$

This means either  $x \in A$  or  $x \in B \cap C,$  and if  $x \in B \cap C$  then  $x \in B$  and  $x \in C$  by definition of intersects.

For the first case,  $x \in A,$   $x$  would also be an element in  $A \cup B$  since it is an element in  $A.$  For the second case,  $x \in B \cap C,$   $x$  would also be an element in  $A \cup B$  since  $x \in B.$

$\therefore$   $A \cup (B \cap C) \subseteq A \cup B.$

Good.

2. Suppose that  $a, b, c \in \mathbb{R}.$  If  $c < 0$  and  $a < b,$  then  $a \cdot c > b \cdot c.$


Proof: Take  $c < 0.$  We can use the Comparison Addition Principle to add  $-c$  to both sides, giving us  $0 < -c.$  Now, we can take  $a < b$  and use the Comparison Multiplication Principle to get  $-ac < -bc.$  Add the quantity  $(ac + bc)$  to both sides (CAP) and you have  $bc < ac,$  which is the same as  $ac > bc.$   $\square$

Excellent!

3. Let  $\mathbb{R}^+ = \{x \mid x \in \mathbb{R} \text{ and } x > 0\}$ . For each  $x \in \mathbb{R}^+$ , let  $A_x = [0, x)$ .


a) What is  $\bigcap_{x \in \{1, 2, 3\}} A_x$ ? *common to all*

$[0, 1)$



b) What is  $\bigcup_{x \in \{1, 2, 3\}} A_x$ ?

$[0, 3)$



c) What is  $\bigcap_{x \in \mathbb{R}^+} A_x$ ?

*there is no least  $\mathbb{R}^+$  so the only intersection is  $\emptyset$  because zero is always included in  $[0, x)$*

$\{0\}$

d) What is  $\bigcup_{x \in \mathbb{R}^+} A_x$ ?

*$\mathbb{R}$  includes all positive reals so any real greater than zero is included*

$[0, \infty)$

*Great!*

4. Suppose  $I$  is a set and for each  $i \in I$ ,  $A_i$  and  $B_i$  are sets, and that there is some  $i \in I$  for which  $A_i \subseteq B_i$ .

a) Is it true that  $\bigcup_{i \in I} A_i \subseteq \bigcup_{i \in I} B_i$ ? Support your answer.

Nope. Suppose  $I = \{1, 2\}$ , with  $A_1 = \{1\}$ ,  $A_2 = \mathbb{R}$ ,  
 $B_1 = \{1\}$ ,  $B_2 = \{2\}$ .

Then  $\bigcup_{i \in I} A_i = \mathbb{R}$  and  $\bigcup_{i \in I} B_i = \{1, 2\}$ , so  $\bigcup_{i \in I} A_i \not\subseteq \bigcup_{i \in I} B_i$ .

But there is some  $i \in I$ , namely 1, for which  $A_1 \subseteq B_1$ .

b) Is it true that  $\bigcap_{i \in I} A_i \subseteq \bigcap_{i \in I} B_i$ ? Support your answer.

Nope. Suppose  $I = \{1, 2\}$ , with  $A_1 = \{1\}$ ,  $A_2 = \{2\}$ ,  
 $B_1 = \{1\}$ ,  $B_2 = \emptyset$ .

So there is some  $i \in I$ , specifically 1, for which  $A_1 \subseteq B_1$ .  
 But  $\bigcap_{i \in I} A_i = \{1, 2\}$  and  $\bigcap_{i \in I} B_i = \emptyset$ , so  $\bigcap_{i \in I} A_i \not\subseteq \bigcap_{i \in I} B_i$ .

5.  $\forall x \in \mathbb{R}, -|x| \leq x \leq |x|$ .

Case 1:  $x \geq 0$ , so by definition  $|x| = x$ , and we can also say  $x \leq |x|$ .

Adding  $-x$  to both sides of  $x \geq 0$  gives us  $0 \geq -x$  or  $-x \leq 0$ .

Then we have  $-|x| = -x \leq 0 \leq x \leq |x|$ .

Case 2:  $x < 0$ , so by definition  $|x| = -x$ , and we can also say  $-x \leq |x|$ .

Adding  $-x$  to both sides of  $x < 0$  gives  $0 < -x$ .

Then we have  $-|x| \leq x < 0 < -x \leq |x|$ .

So in either case we have  $-|x| \leq x \leq |x|$ , as desired.  $\square$

